

Test

Based on Nursing Home Compare - Quality Measures (Q1)

Provider Number	Nursing Home Name	Street
335159	ROSS HEALTH CARE CENTER INC	839 SUFFOLK AVENUE
335837	MARIA REGINA RESIDENCE INC	1725 BRENTWOOD ROAD
335696	GURWIN JEWISH NURSING AND REHABILITATION CENTER	68 HAUPPAUGE ROAD




Test

Based on Nursing Home Compare - Quality Measures (Q1)

City	State	ZIP Code	Phone Number
BRENTWOOD	NY	11717	6312734700
BRENTWOOD	NY	11717	6312734500
COMMACK	NY	11725	6317152000




Test

Based on Nursing Home Compare - Quality Measures (Q1)

Percent of long-stay residents whose need for help with daily activities has increased	Footnote-1
	15.4%
	34.4%
	8.1%




Test

Based on Nursing Home Compare - Quality Measures (Q1)

Percent of long-stay residents who self-report moderate to severe pain		Footnote-2
	0.8%	
	8%	
	5%	




Test

Based on Nursing Home Compare - Quality Measures (Q1)

Percent of long-stay high-risk residents with pressure ulcers	Footnote-3
 8%	
 3.1%	
 3.1%	

Test

Based on Nursing Home Compare - Quality Measures (Q1)

Percent of long-stay residents who lose too much weight	Footnote-4
 2.6%	
 5.8%	
 7.8%	


Test

Based on Nursing Home Compare - Quality Measures (Q1)

Percent of long-stay low-risk residents who lose control of their bowels or bladder	Footnote-5
<div></div>	37%
<div></div>	54.7%
<div></div>	67.4%

Test

Based on Nursing Home Compare - Quality Measures (Q1)



Test

Based on Nursing Home Compare - Quality Measures (Q1)

Footnote-6	Percent of long-stay residents with a urinary tract infection
	<div><div></div></div> 2.6%
	<div><div></div></div> 8.1%
	<div><div></div></div> 7.6%

Test

Based on Nursing Home Compare - Quality Measures (Q1)

Footnote-7	Percent of long-stay residents who have depressive symptoms
	<div></div> 29.1%
	<div></div> 0.6%
	<div></div> 0.5%




Test

Based on Nursing Home Compare - Quality Measures (Q1)

Footnote-8	Percent of long-stay residents who were physically restrained
	0.9%
	2.3%
	3.1%

Test

Based on Nursing Home Compare - Quality Measures (Q1)

Footnote-9	Percent of long-stay residents experiencing one or more falls with major injury	
		2.6%
		2.9%
		1.6%


Test

Based on Nursing Home Compare - Quality Measures (Q1)

Footnote-10

Test

Based on Nursing Home Compare - Quality Measures (Q1)



Test

Based on Nursing Home Compare - Quality Measures (Q1)

Footnote-11

Test

Based on Nursing Home Compare - Quality Measures (Q1)

Percent of long-stay residents assessed and given, appropriately, the pneumococcal vaccine


94.8%

99.4%

99%


Test

Based on Nursing Home Compare - Quality Measures (Q1)



Test

Based on Nursing Home Compare - Quality Measures (Q1)



Test

Based on Nursing Home Compare - Quality Measures (Q1)

Footnote-14

Test

Based on Nursing Home Compare - Quality Measures (Q1)

Percent of short-stay residents assessed and given, appropriately, the seasonal influenza vaccine




Test

Based on Nursing Home Compare - Quality Measures (Q1)

Footnote-15


Test

Based on Nursing Home Compare - Quality Measures (Q1)



Test

Based on Nursing Home Compare - Quality Measures (Q1)



Test

Based on Nursing Home Compare - Quality Measures (Q1)

Footnote-17	Percent of long-stay residents who received an antipsychotic medication
	<div></div> 26.8%
	<div></div> 9.9%
	<div></div> 12.4%

Test

Based on Nursing Home Compare - Quality Measures (Q1)

Footnote-18